

Table S9: Projections of simulations onto 2D-surface using two PC1 modes from unperturbed unbound and bound states. Unbound (green) and bound (brown) states are plotted on the same surface for comparison. Index indicates the residue being held rigid.




Table S9: Projections of simulations onto 2D-surface using two PC1 modes from unperturbed unbound and bound states. Unbound (green) and bound (brown) states are plotted on the same surface for comparison. Index indicates the residue being held rigid.




Table S9: Projections of simulations onto 2D-surface using two PC1 modes from unperturbed unbound and bound states. Unbound (green) and bound (brown) states are plotted on the same surface for comparison. Index indicates the residue being held rigid.




Table S9: Projections of simulations onto 2D-surface using two PC1 modes from unperturbed unbound and bound states. Unbound (green) and bound (brown) states are plotted on the same surface for comparison. Index indicates the residue being held rigid.




Table S9: Projections of simulations onto 2D-surface using two PC1 modes from unperturbed unbound and bound states. Unbound (green) and bound (brown) states are plotted on the same surface for comparison. Index indicates the residue being held rigid.




Table S9: Projections of simulations onto 2D-surface using two PC1 modes from unperturbed unbound and bound states. Unbound (green) and bound (brown) states are plotted on the same surface for comparison. Index indicates the residue being held rigid.




Table S9: Projections of simulations onto 2D-surface using two PC1 modes from unperturbed unbound and bound states. Unbound (green) and bound (brown) states are plotted on the same surface for comparison. Index indicates the residue being held rigid.

